

ON ELLIPTICITY AND GAUGE INVARIANCE IN EUCLIDEAN QUANTUM GRAVITY

Ivan G. Avramidi¹ and Giampiero Esposito²

¹*Department of Mathematics, The University of Iowa,
14 MacLean Hall, Iowa City, IA 52242-1419, USA*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Napoli,
Mostra d'Oltremare Padiglione 20, 80125 Napoli, Italy*

Abstract. Invariance principles determine many key properties in quantum field theory, including, in particular, the appropriate form of the boundary conditions. A crucial consistency check is the proof that the resulting boundary-value problem is strongly elliptic. In Euclidean quantum gravity, the appropriate principle seems to be the invariance of boundary conditions under infinitesimal diffeomorphisms on metric perturbations, and hence their BRST invariance. However, if the operator on metric perturbations is then chosen to be of Laplace type, the boundary-value problem for the quantized gravitational field fails to be strongly elliptic. A detailed proof is presented, and the corresponding open problems are discussed.

Paper prepared for the Conference *Trends in Mathematical Physics*, The University of Tennessee, Knoxville, October 14-17, 1998.

A very important property that emerges from the work in the recent literature^{1–4} is that the request of complete gauge invariance of the boundary conditions is sufficient to determine the most general form of local boundary-value problem for operators of Laplace type. It is therefore quite crucial to understand whether this scheme can be applied to Euclidean quantum gravity in such a way that the ellipticity of the theory is not spoiled. Indeed, despite the lack of a rigorous mathematical framework for a quantization of the gravitational field in the Euclidean regime, several good motivations exist for continuing such an investigation.² In particular, we would like to stress what follows.

(i) Quantum gravity via Euclidean path integrals makes it possible to deal with a framework where, quite naturally, partition functions are defined. This is, in turn, crucial if one wants to “combine” quantum theory, whose predictions are of statistical nature, with general relativity.

(ii) Spectral geometry plays a crucial role if one tries to get a thorough understanding of the one-loop semiclassical approximation. In a space-time approach,² such an approximation provides the “bridge” between the classical world and the as yet unknown full quantum theory (via path integrals). At some stage, any alternative approach to quantum gravity should be able to make contact with what one knows from a perturbative evaluation of transition amplitudes.

(iii) So much has been learned from the heat-kernel approach to index theory and to the theory of eigenvalues in Riemannian geometry, that any new result may have a non-trivial impact on Euclidean quantum gravity. If one is interested in the most fundamental structures, it is encouraging to realize that a framework exists where some problems become well posed. From this point of view, one has to go ahead as long as possible with Euclidean methods, and the problem is open of whether one should regard the elliptic boundary-value problems as the most fundamental tool.

In our investigation, following Refs. 3,4, we first have to describe the vector bundle V over the compact Riemannian manifold (M, g) , the ghost operator and the gauge-field operator with their leading symbols, the projector and the endomorphisms occurring in

the boundary operator. Indeed, in Euclidean quantum gravity, V is the vector bundle of symmetric rank-two tensor fields over M , with bundle metric

$$E^{ab\ cd} = g^{a(c} g^{d)b} + \alpha g^{ab} g^{cd}, \quad (1)$$

which can be inverted whenever $\alpha \neq -\frac{1}{m}$. The generator $\mathcal{R}$ of infinitesimal gauge transformations is the Lie derivative of the symmetric rank-two tensor field φ along the vector field hereafter denoted by ε , i.e.

$$\left(\mathcal{R}\varepsilon\right)_{ab} \equiv (L_\varepsilon\varphi)_{ab} = \sqrt{2} \nabla_{(a} \varepsilon_{b)}, \quad (2)$$

where $\sqrt{2}$ is a normalization factor. The corresponding adjoint generator, say $\overline{\mathcal{R}}$, has an action defined by

$$\left(\overline{\mathcal{R}}\varphi\right)_a \equiv -\sqrt{2} E_a^{\ bc} \nabla_b \varphi_{cd}. \quad (3)$$

Thus, the ghost operator $L \equiv \overline{\mathcal{R}}\mathcal{R}$ turns out to have the leading symbol³

$$\sigma_L(L; \xi) \varepsilon_a = 2E_a^{\ bc} \xi_b \xi_c \varepsilon_d = \left[\delta_a^b |\xi|^2 + (1 + 2\alpha) \xi_a \xi^b \right] \varepsilon_b. \quad (4)$$

To turn L into an operator of Laplace type, which is part of our programme, it is therefore necessary to set $\alpha = -\frac{1}{2}$. To obtain a positive-definite leading symbol for L , it would instead be enough to require that $\alpha > -1$. Moreover, the leading symbol of the operator $H \equiv \mathcal{R}\overline{\mathcal{R}}$ reads³

$$\sigma_L(H; \xi) \varphi_{ab} = 2\xi_{(a} E_{b)}^{\ cde} \xi_c \varphi_{de} = \left[2\xi_{(a} \delta_{b)}^{(c} \xi^{d)} + 2\alpha \xi_a \xi_b g^{cd} \right] \varphi_{cd}. \quad (5)$$

The gauge-invariant operator Δ resulting from the Einstein–Hilbert action is well known to have leading symbol³

$$\begin{aligned} \sigma_L(\Delta; \xi) \varphi_{ab} = & \left[\delta_{(a}^{(c} \delta_{b)}^{d)} |\xi|^2 + \xi_a \xi_b g^{cd} \right. \\ & \left. - 2\xi_{(a} \delta_{b)}^{(c} \xi^{d)} + \frac{(1 + 2\alpha)}{(1 + m\alpha)} g_{ab} (\xi^c \xi^d - g^{cd}) \right] \varphi_{cd}. \end{aligned} \quad (6)$$

By virtue of Eqs. (5) and (6), the gauge-field operator $F \equiv \Delta + H$ is of Laplace type only if $\alpha = -\frac{1}{2}$, just as for the ghost. Hereafter, we shall instead consider for a while an operator of Laplace type, say P , acting on symmetric rank-two tensor fields with fibre metric (1) for arbitrary α .

We are interested in self-adjoint projection operators $\Pi_{ab}{}^{cd}$ with respect to the bundle metric (1). On denoting by N^a the inward-pointing normal vector field to the boundary of M , and by $q_a{}^b \equiv \delta_a{}^b - N_a N^b$ the tensor field which projects vectors onto ∂M , one obtains the desired projector in the form³

$$\Pi_{ab}{}^{cd} \equiv q_{(a}{}^c q_{b)}{}^d - \frac{\alpha}{(\alpha + 1)} N_a N_b q^{cd}. \quad (7)$$

In the applications, it will be useful to bear in mind that the fibre trace of $\Pi_{ab}{}^{cd}$ is equal to $\frac{m}{2}(m - 1)$. The local boundary conditions for our operator of Laplace type can be originally written in the form

$$\left[\Pi_{ab}{}^{cd} \varphi_{cd} \right]_{\partial M} = 0, \quad (8)$$

$$\left[E^{ab}{}^{cd} \nabla_b \varphi_{cd} \right]_{\partial M} = 0. \quad (9)$$

Equation (9) involves both normal and tangential derivatives of φ , so that (8) and (9) are eventually re-expressed as

$$\begin{pmatrix} \Pi & 0 \\ \Lambda & \mathbb{I} - \Pi \end{pmatrix} \begin{pmatrix} [\varphi]_{\partial M} \\ [\nabla_N \varphi]_{\partial M} \end{pmatrix} = 0, \quad (10)$$

with

$$\Lambda \equiv (\mathbb{I} - \Pi) \left[\frac{1}{2} \left(\Gamma^i \widehat{\nabla}_i + \widehat{\nabla}_i \Gamma^i \right) + S \right] (\mathbb{I} - \Pi). \quad (11)$$

Here, in particular,³

$$\Gamma_{ab}{}^{cd} \equiv -\frac{1}{(1 + \alpha)} N_a N_b e^{i(c} N^{d)} + N_{(a} e_{b)}^i N^c N^d, \quad (12)$$

where e_a^i is a local tangent frame on ∂M , and S is an endomorphism. Thus, on denoting by ζ_i the cotangent vectors on the boundary: $\zeta_j \in T^*(\partial M)$, the matrix $T \equiv \Gamma^j \zeta_j$ occurring

in the condition for strong ellipticity³ of the boundary-value problem (P, B_P) reads, in our case,

$$T = -\frac{1}{(1 + \alpha)} p_1 + p_2, \quad (13)$$

where, having defined $\zeta_a \equiv e_a^i \zeta_i$, one has³

$$(p_1)_{ab}{}^{cd} \equiv N_a N_b \zeta^{(c} N^{d)}, \quad (14)$$

$$(p_2)_{ab}{}^{cd} \equiv N_{(a} \zeta_{b)} N^c N^d. \quad (15)$$

The matrix T is annihilated by the projector Π on the left and on the right, as is necessary to be consistent with the desired scheme. Moreover, on studying the products of the matrices p_1 and p_2 , one is led to consider further projectors given by^{3,4}

$$\rho_{ab}{}^{cd} \equiv \frac{2}{|\zeta|^2} N_{(a} \zeta_{b)} N^{(c} \zeta^{d)}, \quad (16)$$

$$p_{ab}{}^{cd} \equiv N_a N_b N^c N^d. \quad (17)$$

One then finds

$$p_1 p_2 = \frac{1}{2} |\zeta|^2 p, \quad (18)$$

$$p_2 p_1 = \frac{1}{2} |\zeta|^2 \rho, \quad (19)$$

where the matrices p_1 and p_2 are nilpotent: $(p_1)^2 = (p_2)^2 = 0$, whilst ρ and p are mutually orthogonal: $p\rho = \rho p = 0$. Thus, on defining

$$\tau \equiv \frac{1}{\sqrt{2(1 + \alpha)}} |\zeta|, \quad (20)$$

the square of the matrix T takes the simple and elegant form

$$T^2 = -\tau^2 (p + \rho). \quad (21)$$

General formulae for even and odd powers of T are also useful, i.e.

$$T^{2n} = (i\tau)^{2n} (p + \rho), \quad (22)$$

$$T^{2n+1} = (i\tau)^{2n}T. \quad (23)$$

The nilpotent matrices p_1 and p_2 have vanishing trace, whereas the projectors p and ρ have unit trace. Hence one finds

$$\text{tr } T^{2n} = 2(i\tau)^{2n}, \quad (24)$$

$$\text{tr } T^{2n+1} = \text{tr } T = 0, \quad (25)$$

which implies, in turn, that for any function, say f , analytic in the region $|z| \leq \tau$, one has^{3,4}

$$\begin{aligned} f(T) &= f(0)[\mathbb{I} - p - \rho] + \frac{1}{2} \left[f(i\tau) + f(-i\tau) \right] (p + \rho) \\ &+ \frac{1}{2i\zeta} \left[f(i\tau) - f(-i\tau) \right] T, \end{aligned} \quad (26)$$

$$\text{tr } f(T) = \left[\frac{m(m+1)}{2} - 2 \right] f(0) + f(i\tau) + f(-i\tau). \quad (27)$$

Thus, the matrix T has eigenvalues $0, i\tau$ and $-i\tau$, and the matrix T^2 , for non-vanishing ζ_j , has eigenvalues 0 and $-\tau^2 = -\frac{|\zeta|^2}{2(1+\alpha)}$. In other words, for $|\zeta| \neq 0$, the matrix $\left(T^2 + |\zeta|^2 \mathbb{I} \right)$ is positive-definite, so that the boundary-value problem (P, B_P) is strongly elliptic with respect to $\mathbf{C} - \mathbf{R}_+$, if and only if³

$$-\frac{1}{2(1+\alpha)} + 1 > 0, \quad (28)$$

which implies $\alpha > -\frac{1}{2}$. But we know from Eqs. (5) and (6) that, in one-loop Euclidean quantum gravity, the gauge-field operator can be of Laplace type if and only if $\alpha = -\frac{1}{2}$. Thus, we have proved, as a corollary, that the gauge-invariant boundary conditions (8) and (9) make it impossible to achieve strong ellipticity with respect to $\mathbf{C} - \mathbf{R}_+$ in Euclidean quantum gravity.^{3,4}

To appreciate which non-trivial properties result from the lack of strong ellipticity we are now going to prove that the heat-kernel diagonal, although well defined, has a non-integrable behaviour when the geodesic distance from the boundary, say r , tends to zero

(unlike the smooth behaviour which is obtained when strong ellipticity is not violated). Indeed, from the work in Ref. 3, one knows that the fibre trace of the zeroth-order heat-kernel diagonal reads

$$\mathrm{tr}_V U_0(x, x; t) = (4\pi t)^{-m/2} \left[C_0 + C_1 e^{-r^2/t} + J(r/\sqrt{t}) \right], \quad (29)$$

where

$$C_0 \equiv \dim V = \frac{m(m+1)}{2}, \quad (30)$$

$$C_1 \equiv \mathrm{tr}_V (\mathbb{I} - 2\Pi) = -\frac{m(m-3)}{2}, \quad (31)$$

$$\begin{aligned} J(z) &\equiv -2 \int_{\mathbf{R}^{m-1}} \frac{d\zeta_1 \dots d\zeta_{m-1}}{\pi^{(m-1)/2}} \\ &\int_{\gamma} \frac{d\omega}{\sqrt{\pi}} e^{-|\zeta|^2 - \omega^2 + 2i\omega z} \mathrm{tr}_V \Gamma^j \zeta_j (\omega \mathbb{I} + \Gamma^k \zeta_k)^{-1}. \end{aligned} \quad (32)$$

By virtue of (27), with $\tau = |\zeta|$ for $\alpha = -\frac{1}{2}$, one finds

$$J(z) = -4 \int_{\mathbf{R}^{m-1}} \frac{d\zeta}{\pi^{(m-1)/2}} \int_{\gamma} \frac{d\omega}{\sqrt{\pi}} e^{-|\zeta|^2 - \omega^2 + 2i\omega z} \frac{|\zeta|^2}{\omega^2 + |\zeta|^2}, \quad (33)$$

where the contour γ comes from $-\infty + i\varepsilon$, goes around the pole $\omega = i|\zeta|$ in the clockwise direction and then goes to $+\infty + i\varepsilon$. The integral with respect to ω is evaluated with the help of the well known formula

$$\int_{\gamma} f(\omega) d\omega = -2\pi i \left[\mathrm{Res} f(\omega) \right]_{\omega=i|\zeta|} + \int_{-\infty}^{\infty} f(\omega) d\omega, \quad (34)$$

and the integrals with respect to $\zeta_1, \dots, \zeta_{m-1}$ (see (32)) can be reduced to Gaussian integrals. This method leads to the result³

$$J(z) = 2(m-1)z^{-m} \Gamma(m/2, z^2), \quad (35)$$

where we have used the incomplete Γ -function $\Gamma(a, x) \equiv \int_x^\infty u^{a-1} e^{-u} du$. The function J is singular as $z \rightarrow 0$, in that

$$J(z) \sim 2(m-1)\Gamma(m/2)z^{-m}, \quad (36)$$

whereas, as $z \rightarrow \infty$, it is exponentially damped, i.e.

$$J(z) \sim 2(m-1)z^{-2}e^{-z^2}, \quad (37)$$

as can be seen by using the identity $\Gamma(a, x) = x^{a-1}e^{-x} + (a-1)\Gamma(a-1, x)$. The singularity at $z = 0$ results from the pole at $\omega = i|\zeta|$ and provides a peculiar difference with respect to the strongly elliptic case, where all poles lie on the positive imaginary line with $\text{Im}(\omega) < i|\zeta|$. The fibre trace (29) is now re-expressed as³

$$\text{tr}_V U_0(x, x; t) = (4\pi t)^{-m/2} \left[C_0 + C_1 e^{-r^2/t} + 2(m-1)(r/\sqrt{t})^{-m} \Gamma(m/2, r^2/t) \right]. \quad (38)$$

One can see that the first term in (38) yields the familiar interior contribution, upon integration over a compact manifold. Moreover, the boundary terms in (38) are exponentially small as $t \rightarrow 0^+$, if r is fixed. By contrast, on fixing t , one finds, as $r \rightarrow 0$,

$$\text{tr}_V U_0(x, x; t) \sim (4\pi)^{-m/2} 2(m-1)\Gamma(m/2)r^{-m}. \quad (39)$$

Remarkably, such a limiting behaviour is independent of t and is not integrable with respect to r as one approaches the boundary, i.e. as $r \rightarrow 0$.³ This is an enlightening example of the undesirable properties resulting from the lack of strong ellipticity.

Our work, relying on Refs. 3,4, has proved that, upon studying Euclidean quantum gravity at one-loop level on compact Riemannian manifolds with smooth boundary, if one tries to have a gauge-field operator and ghost operator both of Laplace type, with local boundary operator (see (10) and (11)) and completely gauge-invariant boundary conditions, the ellipticity of the theory is lost in that the boundary-value problem for metric perturbations fails to be strongly elliptic with respect to $\mathbf{C} - \mathbf{R}_+$. Since the properties of ellipticity on the one hand, and invariance under infinitesimal diffeomorphisms (on metric perturbations) on the other hand, are both at the heart of the Euclidean approach

to quantum gravity, the proof of their incompatibility appears to be a matter of serious concern. At least three alternatives deserve further investigation. They are as follows.

- (i) The use of smearing functions (e.g. $f : r \rightarrow f(r) = r^m$) in defining the functional trace of e^{-tF} , with F the gauge-field operator $\Delta + H$.
- (ii) The attempt to prove strong ellipticity when the gauge-field operator is non-minimal.
- (iii) The attempt to recover strong ellipticity upon studying non-local boundary conditions, with the help of the current progress in the functional calculus of pseudo-differential boundary problems.^{5,6}

The latter line of investigation, although quite difficult at the technical level, might lead to a deeper vision,⁶ and finds its motivation not only in the recent mathematical progress,⁵ but also in well known properties of quantum field theory, e.g. the non-local nature of projectors on the gauge-invariant sub-space of the configuration space.³ There is therefore some intriguing evidence that the most fundamental problems in spectral geometry and quantum gravity are still waiting for a proper solution.

References

- [1] A. O. Barvinsky, *Phys. Lett.*, **B195** (1987) 344.
- [2] G. Esposito, A. Yu. Kamenshchik and G. Pollifrone, *Euclidean Quantum Gravity on Manifolds with Boundary*, Fundamental Theories of Physics, Volume 85 (Kluwer, Dordrecht, 1997).
- [3] I. G. Avramidi and G. Esposito, *Gauge Theories on Manifolds with Boundary* (hep-th 9710048, to appear in *Commun. Math. Phys.*).
- [4] I. G. Avramidi and G. Esposito, *Class. Quantum Grav.*, **15** (1998) 1141.
- [5] G. Grubb, *Functional Calculus of Pseudo-Differential Boundary Problems*, Progress in Mathematics, Volume 65 (Birkhäuser, Boston, 1996).
- [6] G. Esposito, *Non-Local Boundary Conditions in Euclidean Quantum Gravity* (gr-qc 9806057).